



Planta Medica

Journal of Medicinal Plant and Natural Product Research

11

Volume 78
August 2012
Page 1027-1310

8th Joint Meeting of AFERP, ASP, GA, PSE & SIF

28th July–1st August 2012, New York City, USA

Abstracts

Chairmen:
Guy T. Carter
Edward J. Kennelly

Issue Editor:
Mark O'Neil-Johnson



Cistus ladanifer as a source of phenolic compounds with antifungal activity

Barros L^{1,2}, Dueñas M², Alves CT³, Silva S³, Henriques M³, Santos-Buelga C², Ferreira ICFR¹

¹CIMO-ESA Polytechnic Institute of Bragança, Portugal; ²GIP Faculty of Pharmacy, University of Salamanca, Spain; ³IBB Centre of Biological Engineering, University of Minho, Braga, Portugal

A screening of the antifungal potential of phenolic extract of *Cistus ladanifer* from Northeast Portugal, against *Candida* species was performed. The extract was characterized by HPLC-DAD-ESI/MS. Phenolic acids and derivatives, ellagic acid derivatives and flavonoids, such as catechins, flavonols and flavones, were found in the sample. The most abundant group was ellagic acid derivatives in which punicalagin gallate, a derivative of punicalagin attached to gallic acid, was found in highest amount. These compounds could be related to the strong inhibition of *C. albicans*, *C. glabrata* and *C. parapsilosis* growth. Moreover, the best antifungal activity was against *C. glabrata*, where the studied extract was able to cause at least 3 Log of reduction at concentrations below 50 µg/mL and a total growth inhibition at concentrations above 625 µg/mL.

CISTUS LADANIFER AS A SOURCE OF PHENOLIC COMPOUNDS WITH ANTIFUNGAL ACTIVITY

Lillian Barros^{1,2}, Montserrat Dueñas², Carlos Tiago Alves³, Sónia Silva³, Mariana Henriques³, Celestino Santos-Buelga², and Isabel C.F.R. Ferreira¹. ¹CIMO-ESA Polytechnic Institute of Bragança, Portugal, ²GIP Faculty of Pharmacy, University of Salamanca, Spain, ³IBB Centre of Biological Engineering, University of Minho, Braga, Portugal.

A screening of the antifungal potential of phenolic extract of *Cistus ladanifer* from Northeast Portugal, against *Candida* species was performed. The extract was characterized by HPLC-DAD-ESI/MS. Phenolic acids and derivatives, ellagic acid derivatives and flavonoids, such as catechins, flavonols and flavones, were found in the sample. The most abundant group was ellagic acid derivatives in which punicalagin gallate, a derivative of punicalagin attached to gallic acid, was found in highest amount. These compounds could be related to the strong inhibition of *C. albicans*, *C. glabrata* and *C. parapsilosis* growth. Moreover, the best antifungal activity was against *C. glabrata*, where the studied extract was able to cause at least 3 Log of reduction at concentrations below 50 µg/mL and a total growth inhibition at concentrations above 625 µg/mL.

Acknowledgements: PEst-OE/AGR/UI0690/2011, SFRH/BPD/4609/2008 (L. Barros), Ramón y Cajal (M. Dueñas).